



# Effect of Autoclaving on Proximate and Anti-Nutritional Factors of Jackbean (*Canavalia ensiformis* L) D.C. FLOUR

Odedeji, J.O.<sup>1</sup>, Akande, E.A.<sup>2</sup>, Alade O.A.<sup>2</sup> and Ayinde L.A.<sup>1</sup>.

<sup>1</sup>Food Science and Technology Department, Osun State Polytechnic, Iree, Osun State, Nigeria.

<sup>2</sup>Chemical Engineering Department, Ladoko Akintola University of Technology, Ogbomoso, Oyo State, Nigeria.

<sup>1</sup>Statistics Department, Osun State Polytechnic, Iree, Osun State, Nigeria.

E-mail: odedeji2014@gmail.com

Received: July 15, 2020

Accepted for publication: August 17, 2020

Published: November 20, 2020

## Abstract

Jackbean (*Canavalia ensiformis*) is a legume with high nutritive value but with elongated cooking time, hard seed coat and anti-nutritional factors as limiting factors. There is a dearth of information on the impact of different pre-processing methods on the proximate and anti-nutritional factors of Jackbean flour. Therefore, this study is on the influence of autoclaving on proximate and anti-nutritional factors of Jackbean flour. Jackbean procured from the Genetic Resources Unit of International Institute of Tropical Agriculture (IITA), Ibadan, Oyo State Nigeria were subjected to autoclaving at 121 °C for 30, 60, 90, 120 mins, dried to 12 % moisture content and milled into flour while flour from untreated seed served as control. The data generated were subjected to statistical analyses using SAS 2.0 and SPSS 20.0 packages. The results revealed that autoclaving at 121 °C for 120 mins resulted in maximum protein content of 34.97 % and drastic reduction in all anti-nutritional factor most particularly trypsin inhibitor from 27.88 % in raw seed to 17.86 % in autoclaved seed. Autoclaving thus had positive effect on the proximate and anti-nutritional factors of Jackbean.

**Keywords:** Proximate, anti-nutritional, autoclaving, jackbean

## Introduction

Jackbean (*Canavalia ensiformis*) – “Sese nla” (Yoruba), is one of the tropical legumes that do well in Nigeria (Okonkwo and Udedibie, 1991). It is grossly under-utilized in most developing countries of the world most particularly Nigeria. This has really affected the maximum propagation of the crop despite its nutritional qualities. Some people plant the seeds in Nigeria as ornamental plants and in some places, it is believed to be “snake repellent”. Jackbean has potential in product development and consumption, but its prolonged period of processing, and tough seed coat have affected the properties of the flour (Akande et al., 2013). Quality parameters of food products developed from legume flour samples generally affect the functional and pasting properties influenced by the processing conditions employed (Ezeocha and Onwuka, 2010).

Various pre-processing techniques had been employed to improve the processing, nutritional quality, organoleptic acceptability, reduction in the oligosaccharides and other anti-nutritional factors. Some of the commonly used pre-processing techniques included soaking, boiling at high temperatures (in water, alkaline or acidic solutions), sprouting,

roasting, dehulling, microwaving, steam blanching, fermentation and autoclaving (El-Adawy, 2002 and Skulinova *et al.*, 2002). Autoclaving had been exploited to a great extent in food processing and found to be more effective in the reduction of anti-nutritional components of leguminous products (Adegunwa *et al.*, 2012).

Food processing operations not only improves the flavor and palatability of foods but also increase the bioavailability of nutrients by inactivating some anti-nutritional factors and growth inhibitors (Chau *et al.*, 1997). Studies had shown that pre-treatments such as sprouting, soaking/fermentation, autoclaving, blanching or roasting could enhance the quality of seeds and grain products (Ade- Omowaye *et al.*, 2003; Kirbaslar and Erkmn, 2003 and Carmelia *et al.*, 2007). Sprouting of soybean had been reported to increase the ascorbic acid content, riboflavin and niacin (Iwe, 2003).

Reported studies on the beans are majorly on its utilization in animal feed formulations (Udebibie, 1997 and Anyanwu *et al.*, 2011). There is a paucity of information on the effect of autoclaving on the proximate and anti-nutritional factors of Jackbean grown in developing countries, which is responsible for its in-extensive use (Ikegwu *et al.*, 2010). This work therefore seeks to investigate the effect of autoclaving on the proximate and anti-nutritional factors of Jackbean flour.

## Materials and Methods

### *Source of Planting Material and Experimental Location*

The seeds of Jackbean (*Canavalia ensiformis*) with Accession Number: TCe4 were collected from the Genetic Resources Unit of International Institute of Tropical Agriculture (IITA), Ibadan, Oyo State Nigeria. The seeds were planted at the experimental farm of Osun State Polytechnic, Iree, Osun State, Nigeria. Chemicals (Analytical Grade) and equipment were obtained from the Department of Food Science and Engineering, Ladoke Akintola University of Technology, Ogbomosho and Department of Food Technology, Osun State Polytechnic, Iree, Osun State.

### *Preparation of the beans*

The pods of Jackbean were harvested after four month of maturity and sundried. The seeds were carefully removed from the pod by shelling and were winnowed thereafter so as to remove chaff.

### *Production of Flour from Raw and Autoclaved Jackbean*

Matured Jackbean seeds, after four months of planting, were thoroughly cleaned and sorted to remove defective ones. The seeds were thereafter graded according to their sizes. The cleaned seeds were soaked in warm water to soften the hull and ease its removal. The hulls were removed by repeated working between the palms and were removed leaving behind dehulled seeds. The dehulled seeds were thereafter drained and dried to reduce the moisture content and facilitate grinding into flour.

Two hundred grammes (200g) of the seed samples were autoclaved at 151b pressure (121 °C) in distilled water in 1:10 ratio (bean: water) for 30, 60, 90 and 120 minutes. The autoclaved seeds were dehulled and oven dried at 55 °C. The dried seeds were then milled in attrition mill, sieved to 1mm mesh size, and packaged in an air-tight container for further analytical work (Adegunwa *et al.*, 2012; Uche *et al.*, 2014).

### *Effect of Pre-treatments on Proximate Constituents of Jackbean Flour*

#### *Moisture content and dry matter*

The moisture content was determined using the method described by No. 925.10, AOAC (1990). The percentage moisture content and dry matter were calculated as follows:

$$\% \text{ Moisture content} = \frac{(W_2 - W_3)}{(W_2 - W_1)} \times 100$$

$$\% \text{ Dry matter} = \frac{W_2 - W_0}{W_1 - W_0} \times 100\%$$

$W_1$  = Weight of crucible

$W_2$  = Weight of crucible + dry sample

$W_3$  = Weight of crucible + dry sample with constant weight.

#### *Crude protein*

The crude protein content was determined using kjedhal method as described by AOAC, (2000).

$$\%N = \frac{1.401 \times (\text{cm}^3 \text{ HCl sample} - \text{cm}^3 \text{ HCl titre of blank}) \times \text{MHCl}}{S_w}$$

% protein = % N x 6.25

Sw = Sample weight

N = Nitrogen

#### *Ash content*

The ash content was determined using AOAC, (1990) method.

$$\% \text{ Ash content} = \frac{W_2 - W_0}{W_1 - W_0} \times 100\%$$

$W_0$  = Weight of the crucible

$W_1$  = Weight of the crucible + sample before ashing

$W_2$  = Weight of the crucible + sample after ashing

#### *Crude fat*

Five (5) g of the sample was measured into a set of three free thimbles after it has been dried free of moisture (AOAC, 2000) using soxhlet apparatus.

$$\% \text{ Crude fat (ether extract)} = \frac{W_1 - W_2}{S_w} \times 100\%$$

$W_1$  = Weight of the flask

$W_2$  = Weight of the flask and the oil

Sw = Sample Weight

#### *Crude fibre*

Crude fibre was determined through the documented method of AOAC, (2000).

$$\% \text{ Crude fibre} = \frac{W_1 - W_2}{S_w} \times 100\%$$

$W_1$  = Weight before incineration

$W_2$  = Weight after incineration

Sw = Sample Weight

*Effect of Pre-treatment on the Anti-nutritional Factors of the Flour**Phytate content*

The phytate content of the flour was determined using the method of Maga (1982). 2g of each finely ground flour sample was soaked in 20 ml of 0.2N HCl filtered, 0.5 ml filtrate was mixed with 1 ml Ferric ammonium sulphate solution, boiled for 30 min, cooled in ice for 15mins and centrifuged for 15 min. One milliliter of the supernatant was mixed with 1.5 ml of 2.2 – pyridine solution and the absorbance measured in a spectrophotometer at 519nm.

*Tannin content*

Tannin content of the samples were determined using the method of Makkar and Goodchild, (1996).

*Oxalate content*

Titration method described by Day and Underwood (1986) was used to determine the oxalate content.

*Polyphenols content*

The total polyphenols in the sample was determined using pursson blue spectrophotometric method with tannic acid as the standard as described by Oladele *et al.* (2009).

*Trypsin inhibitor*

The modified method of Arntified *et al.* (1985) was used in the determination of trypsin inhibitor which involves the use of spectrophotometer to measure the absorbance of supernatant of centrifuged mixture of 0.5g of the sample with 50 mls of 0.5m NaCl solution. The absorbance for the control was measured. Trypsin inhibitor activity was calculated using the formula:

$$\text{Tul/g} = \frac{1}{w} \times \frac{au - as}{0.01} \times \frac{vf}{va}$$

Where:

- w = Weight of sample
- au = Absorbance of sample at 410nm
- as = Absorbance of control
- vf = Total extract volume
- va = Volume of extract analyzed

*Saponin content*

The solvent extraction gravimetric method described by Nwosu, (2010) was used for the determination of saponin content of the sample.

$$\% \text{ Saponin} = \frac{W_2 - W_1}{\text{Wt of sample}} \times \frac{100}{1}$$

Where:

- W<sub>1</sub> = Weight of evaporating dish
- W<sub>2</sub> = Weight of dish of sample

*Data Analysis*

Data collected were subjected to analysis of variance (ANOVA) using SPSS version 20.0 and SAS 2.0 package. The means were separated by Duncan's multiple range test at 5% level of probability, while Pearson's correlation co-efficient was also done to establish relationship among the preprocessing methods used in this research.

## Results and Discussion

### *Effect of Autoclaving on Proximate Composition of Jackbean Flour*

The result of the effect of autoclaving on proximate composition of Jackbean flour is presented in Table 1.0. Sample T1 showed higher significant effect ( $P < 0.05$ ) and mean of  $8.73 + 0.46$  for moisture content compared to other samples and the raw sample. The raw sample and sample T4 showed no significant difference but lower mean value of  $8.33 + 0.23$  for moisture content. Moisture content was observed to be increasing among the pre-processed samples over the period of autoclaving except for sample T4 when there is drop in the moisture content to  $8.33 + 0.11$ . The moisture content of the samples was within the recommended range of 10-15 for dry products (Adegunwa *et al.*, 2012).

The protein content of samples T3 and T4 displayed higher significant ( $P < 0.05$ ) effect and mean values of  $34.91 + 0.35$  and  $34.97 + 0.15$  respectively compared to the raw sample. The values ranged between  $34.97 + 0.15$  and  $32.02 + 0.08$  while the raw sample recorded the least value. The protein content was increasing over the autoclaving period. This might be due to the elimination of some anti-nutritional factors. The values recorded for protein were higher compared to the value reported for autoclaved Beniseed (Adegunwa *et al.*, 2012). This indicated that the sample could be a cheap source of plant protein.

The mean values of  $3.80 + 0.10$  and  $3.72 + 0.08$  was recorded for samples T1 and the untreated sample for fat content compared to other samples. There was decrease in fat content as the autoclaving period increased from  $3.72 + 0.08$  to  $3.43 + 0.06$ . Sample T4 recorded the least value. This might be due to evaporative loss during the process of drying. The ash content showed a significant effect ( $P < 0.05$ ) and mean value of  $4.43 + 0.11$  for the untreated sample while other samples present no significant effect and mean values ranging from  $4.00 + 0.00$  to  $4.08 + 0.10$ . There was reduction in ash content over the autoclaving period. The ash content is relatively lower compared to the value recorded for benniseed (Adegunwa *et al.*, 2012). Similar significant effect ( $P < 0.05$ ) but mean values of  $5.01 + 0.34$  and  $4.99 + 0.33$  were recorded for samples T0 and T1 which was followed by samples T2 and T3 with mean values of  $4.71 + 0.11$  and  $4.63 + 0.06$  respectively while sample T4 recorded the last value of  $4.52 + 0.07$ . The crude fibre is the amount of indigestible sugar present in a food sample. The values were however higher than the value reported for cooked African Yam Bean (*Spherostylis sternocarpa*) seed as reported by Nwosu *et al.* (2011).

**Table 1: Effect of Autoclaving on Proximate Composition of Jackbean Flour**

Parameters (%)	T0	T1	T2	T3	T4
Moisture Content	$8.33 + 0.23c$	$8.73 + 0.46a$	$8.60 + 0.40b$	$8.47 + 0.23ab$	$8.33 + 0.11c$
Protein	$32.02 + 0.08c$	$31.42 + 0.02d$	$33.13 + 0.11b$	$34.91 + 0.35a$	$34.97 + 0.15a$
Fat	$3.80 + 0.10a$	$3.72 + 0.08a$	$3.56 + 0.06b$	$3.48 + 0.03bc$	$3.43 + 0.06c$
Ash	$4.43 + 0.11a$	$4.08 + 0.10b$	$4.01 + 0.01b$	$4.00 + 0.00b$	$4.00 + 0.00b$
Crude fibre	$5.01 + 0.34a$	$4.99 + 0.33a$	$4.71 + 0.11ab$	$4.63 + 0.06ab$	$4.52 + 0.07b$
Carbohydrate	$46.42 + 0.43b$	$47.02 + 0.22a$	$45.99 + 0.21b$	$44.51 + 0.40c$	$44.62 + 0.10c$

*Values are means of triplicate determinations. Means with the same superscript are not significantly different ( $P < 0.05$ ) along the column*

*T0 = Raw Jackbean Flour*

*T1 = Autoclaved jackbean flour at 121 oC for 30 min*

*T2 = Autoclaved jackbean flour at 121 oC for 60 min*

*T3 = Autoclaved jackbean flour at 121 oC for 90 min*

*T4 = Autoclaved jackbean flour at 121 oC for 120 min*

Higher significant effect ( $P < 0.05$ ) and mean of  $47.02 + 0.22$  was displayed by sample T1 for carbohydrate content compared to other pre-processed samples and the untreated sample. The value was however found to decrease from  $47.02$

+ 0.22 to 44.62 + 0.10 over the autoclaving period. Samples T3 and T4 showed the lowest means of 44.51 + 0.40 and 44.62 + 0.10 but similar significant effect for carbohydrate.

The improvement in the proximate constituents of the sample due to the effect of autoclaving supported the views of Adegunwa *et al.* (2012); Doss *et al.* (2011) and Nwosu *et al.* (2011) on biochemical composition, anti-nutritional, proximate, and nutritional values of beniseed, underutilized legume and African yam bean respectively. The reduction in carbohydrate and lipid content in this sample is beneficial for patients of diabetes mellitus, cardiovascular disease and hypercholesterolemia (Myrene, 2013). The carbohydrate content was relatively low compare to the value reported for autoclaved field bean by Myrene (2013).

#### *Effect of Autoclaving on Anti-nutritional Factors of Jackbean Flour*

The results of the effect of autoclaving on anti-nutritional factors of Jackbean flour is shown in Table 2. There was higher significant effect ( $P < 0.05$ ) with a mean value of 1.99 + 0.07 on phenolic compound by raw sample. This was closely followed by sample T1 with a mean value of 1.72 + 0.11 while samples T2 and T3 showed no significant effect but have mean values of 1.47 + 0.06 and 1.42 + 0.02 respectively. Sample T4 recorded the least value of 1.05 + 0.10 for this anti-nutritional factor. There was gradual reduction of phenolic compound as the autoclaving time increased, which corroborate the observations of Udensi *et al.* (2008) they recorded lower value for phenolic compound for *Mucuna utilis* and velvet beans respectively.

There was variation in tannin content among all the samples over the autoclaving times. Higher significant effect ( $P < 0.05$ ) with a mean of 1.99 + 0.07 was recorded by the untreated sample while samples T2 and T3 showed similar significant effect but varied mean values of 1.47 + 0.06 and 1.42 + 0.02 respectively. The least content of 0.73 + 0.06 for tannin was recorded by sample T1. Tannins are astringent and adversely affect feed intake (Microsoft Encarta, 2006). Autoclaving as a form of pre-processing method thus reduced this anti-nutritional factor drastically. The phytate causes reductions in absorption of calcium from the gastrointestinal tract and consequently implicated in the development of ricket (Deshpande and Cheryan, 1984). The value for phytate ranged between 0.52 + 0.04 and 0.14 + 0.03. Higher significant ( $P < 0.05$ ) effect was recorded by sample T4 with a mean value of 0.52 + 0.04 while the raw sample recorded the least value. Autoclaving had an incremental effect on phytate. Higher significant ( $P < 0.05$ ) effect and mean value of 0.12 + 0.01 was recorded for raw sample for oxalate compared to other autoclaved samples.

**Table 2: Effect of Autoclaving on Anti-nutritional Factors of Jackbean Flour**

Parameters	To	T1	T2	T3	T4
Phenolic (mg/g)	1.99 + 0.07a	1.72 + 0.11b	1.47 + 0.06c	1.42 + 0.02c	1.05 + 0.10d
Tannin (mg/g)	1.99 + 0.07a	0.73 + 0.06d	1.47 + 0.06b	1.42 + 0.02b	1.05 + 0.10c
Phytate (mg/g)	0.14 + 0.03d	0.13 + 0.02e	0.17 + 0.01c	0.37 + 0.01b	0.52 + 0.04a
Oxalate (%)	0.12 + 0.01a	0.10 + 0.01b	0.07 + 0.01c	0.06 + 0.01c	0.05 + 0.01c
Saponin (%)	0.75 + 0.05bc	0.91 + 0.07ab	0.89 + 0.02ab	0.79 + 0.02b	0.68 + 0.02c
Trypsin Inhibitor (%)	27.88 + 0.01a	23.38 + 0.01b	20.42 + 0.03c	19.80 + 0.03d	17.86 + 0.16e

Values are means of triplicate determinations. Means with the same superscript are not significantly different at  $P < 0.05$  along the column

To = Raw jackbean flour

T1 = Autoclaved jackbean flour at 121 oC for 30 min

T2 = Autoclaved jackbean flour at 121 oC for 60 min

T3 = Autoclaved jackbean flour at 121 oC for 90 min

T4 = Autoclaved jackbean flour at 121 oC for 120 min

There was reduction of oxalate content as the autoclaving time increased. The value reduced from 0.10 + 0.01 in sample T1 to 0.05 + 0.01 in sample T4 while samples T2, T3 and T4 showed similar significant effect. This reducing effect is in line with the submission of Nwosu (2010) who reported reduction in oxalate content of Asparagus bean. The content of oxalate in

the sample is relatively low compared to other anti-nutritional factors. Significant reduction in saponin was recorded when the sample was autoclaved for 120 min (T4) with a mean value of  $0.68 + 0.01$ . There was a steady reduction in the content of saponin from  $0.91 + 0.01$  in sample T1 to  $0.68 + 0.01$  in sample T4 indicating that autoclaving had a reducing effect on this anti-nutritional factor. There is no significant effect recorded for samples T1 and T2 based on saponin content.

There is drastic reduction of trypsin inhibitor from  $27.88 + 0.01$  in sample T0 to  $17.86 + 0.01$  in sample T4 due to autoclaving. Sample T4 recorded the least value for trypsin inhibitor while highest value of  $27.88 + 0.01$  was recorded in the raw sample. Autoclaving, a thermal treatment, thus had a reducing effect on this heat labile anti-nutritional factor thus corroborating the submission of Myrene, (2013); Adegunwa et al. (2012) on the effect of traditional and thermal processing on nutritional and biochemical, antinutritional and functional properties of field bean and beniseed.

### Conclusion

The study revealed that autoclaving had significant effects on proximate and anti-nutritional factors. Autoclaving at 120 °C for 120 min resulted in significant reduction of anti-nutritional factors especially trypsin inhibitor and increment in protein and carbohydrate contents associated with low moisture. Autoclaving of Jackbean therefore represents a pre-processing technique with positive effect on proximate and anti-nutritional factors. With this treatment properly applied, Jackbean could be crop of great industrial potentials, thus, its maximum propagation should be encouraged.

### Acknowledgements

All the authors whose research works were referenced and cited are hereby acknowledged.

### Conflict of Interests

The authors declare that there is no conflict of interest.

### References

- Adegunwa MO, Adebawale AA and Solano EO (2012). Effect of Thermal Processing on the Biochemical Composition, Anti-nutritional Factors and Functional Properties of Beniseed (*Sesamum Indicum*) Flour. *American Journal of Biochemistry and Molecular Biology* 2(3): 175 – 182.
- Ade-Omowaye BIO, Olajide JO and Oluyomi EO (2003). Pretreatment Sorghum-Cassava Flour as a Substitute for Traditional Nigeria Yam Flour (Elubo) *Plant Foods Human Nutrition*. 58: 1 – 11.
- Akande EA, Oladipo AO and Kelani OS (2013). Effect of Steaming on The Physicochemical Properties and The Cooking Time of Jackbean (*Canavalia ensiformis*). *Journal of Experimental Biology and Agricultural Sciences*. 1: 6.
- Anyanwu DC, Ukaegbu EP and Ogueri C (2011). Nutrient Utilization and Growth Response of *Clarias Ganepienus* Fingerlings Fed Dietary Levels of Jackbean (*Canavalia ensiformis*) Meal. *New York Science Journal*. 4(4): 68-71.
- AOAC (1990). *Official Methods of Analysis*. 15th Edition, Association of Official Analytical Chemist. Washington D.C.
- AOAC (2000). *Association of Official Analytical Chemists. Official Methods of Analysis*. 17th ed. Washington D.C. Pp. 16
- Arntfield SD, Ismopnd MAH and Murray ED (1985). The Fate of Anti-nutritional Factors during the Preparation of Faba Bean Protein Isolate using Micellization Techniques. *Can.inst. Food Science. Innov.* 18: 137 – 143.
- Carmelia, A. S., Beatriz de la C., Marcel B., Richard H. S., Elke A. (2007). Impact of the Roasting Degree of Coffee on the in Vitro Radical Scavenging Capacity and Content of Acrylamide *LWT* 40 : 1849 – 1854. .

Chau CF, Cheung PC and Wong YS (1997). Effect of cooking on content of amino acids and antinutrients in three Chinese indigenous legume seeds. *Journal of Science of Food and Agriculture*. 75: 447-452.

Day RA (Jnr) and Underwood AL (1986). *Quantitative Analysis*. 5th Ed. Prentize – Hall Publication.

Desphande S and Chayan M (1984). Changes in Phytic Acid, Tannins and Trypsin Inhibitor Activity on Soaking of Dry Beans (*Phaseolus vulgaris* L.) *Nutrition*. 27: 371 – 377.

Doss A, Pugalenti M, Vadivel VG, Subhashini G and Anitha-Subash RA (2011). Effects of processing technique on the nutritional composition and antinutrients content of under-utilized food legume *Canavalia ensiformis* L. Dc. *International Food Research Journal*. 18(3): 965-970.

EI – Adawy TA (2002). Nutritional Composition and Anti-nutritional Factors of Chickpeas (*Cicer arietinum* L) Undergoing Different Cooking Methods and Germination. *Plant Food for Human Nutrition*. 57: 83- 97.

Ezeocha VC and Owuka GI (2010). Effect of Processing Methods on the Physicochemical and Nutritional Quality of Maize and soybean based complementary Blends. *Nigerian Food Journal*. 28(2): 210 – 216.

Ikwegwu OJ, Okechukwu PE and Ekumankama EO (2010). Physico-chemical and Pasting Characteristics of Flour and Starch from 'Achi' *Brachystegia eurycoma* Seed. *Journal of Food Technology*. 8(2): 58 – 66.

Iwe MO (2003). *The Science and Technology of Soybean: Chemistry, Nutrition, Processing and Utilization*. Rojoint Communication Services Limited Publisher.

Kirbaslar FG and Erkmén G (2003). Investigation of the Effect of Roasting Temperature on the Nutritive Value of Hazelnuts Plant Foods. *Hum Nutri*. 58: 1 – 10.

Maga JA (1982). Phytates: Its Chemistry, Occurrence, Food Interactions, Nutritional Significance and Methods of Analysis. *Journal. Agric. Food Chem*. 30(1): 1 – 7.

Makkar AOS and Goodchild AV (1996). Qualification of Tannins. A Laboratory Manual Int. Centre for Agriculture. A Laboratory Manual Int. Centre for Agriculture Research in the Dry Areas (ICARDA), Aleppo, Syria, I.V + 25pp.

Microsoft @ Encarta ® 2006 (CD). Redmond WA (2006). Microsoft Corporation

Myrene R D'souza (2013). Effect of Traditional Processing on Nutritional Quality of Field Bean. *Advances in BioResearch*. 4 (3): 29 – 33.

Nwosu JN (2010). Effect of Soaking, Blanching and Cooking on the Anti-nutritional Properties of Asparagus Bean (*Vigna Sesquipedis*) Flour. 1-5.

Nwosu JN, Ahaotu I, Ayozie C, Udeozor LO and Ahaotu NN (2011). The proximate and functional properties of African Yam Bean (*Sphenostylis sternocarpa*) Seeds as affected by processing. 29 (2): 39-48.

Okonkwo JC and Udedibie ABI (1991). Preliminary observation on the yield performance of Jackbean (*Canavalia ensiformis*) and sword bean (*Canavalia gladiata*) in the Guinea Savannah of Nigeria. Paper presented at the 27th Annual Conference of Agriculture of Nigeria, Mina, Nigeria. 1 – 4 September.

Oladele AK, Osundahunsi OF and Adebowale YA (2009). Effect of Processing Techniques on the Nutrients and Antinutrients Contents of Tigernut (*Cypercus esculentus*). *Nigerian Food Journal*. 27(2): 210 – 218.

Skulinova M, Kadlec P, Kaasova J, Dostalora J, Zatopkova M, Mosnedl V and Hrachovivova J (2002). Microwave Treatment and Drying of Germinated Pea. *Zech. Journal of Food Science*. 20: 23 - 30.

Uche SN, Charity UN, Abbas O, Aliyu M, Francis CB and Oche O (2014). Proximate, antinutrients and mineral composition of raw and processed (boiled and roasted) *Sphenostylis sternocarpa* seeds from Southern Kaduna North-West Nigeria. *ISRN. Nutrition*.

Udebibie ABI (1997). Jackbean (*Canavalia ensiformis*) Meal in Poultry Diets. In *Sustainable Rural Poultry Production in Africa*. E.B. Sonaiya (ed.), pp. 97-107

Udensi EA, Onwuka GI and Okoli EG (2008). Effect of processing on the levels of some antinutritional factors in *Mucuna utilis*. *Plant Products Research Journal*. 8(1): 1-6.